

IZA DP No. 9224

**Do More of Those in Misery Suffer from Poverty,
Unemployment or Mental Illness?**

Sarah Flèche
Richard Layard

July 2015

Do More of Those in Misery Suffer from Poverty, Unemployment or Mental Illness?

Sarah Flèche

CEP, London School of Economics

Richard Layard

*CEP, London School of Economics
and IZA*

Discussion Paper No. 9224
July 2015

IZA

P.O. Box 7240
53072 Bonn
Germany

Phone: +49-228-3894-0
Fax: +49-228-3894-180
E-mail: iza@iza.org

Any opinions expressed here are those of the author(s) and not those of IZA. Research published in this series may include views on policy, but the institute itself takes no institutional policy positions. The IZA research network is committed to the IZA Guiding Principles of Research Integrity.

The Institute for the Study of Labor (IZA) in Bonn is a local and virtual international research center and a place of communication between science, politics and business. IZA is an independent nonprofit organization supported by Deutsche Post Foundation. The center is associated with the University of Bonn and offers a stimulating research environment through its international network, workshops and conferences, data service, project support, research visits and doctoral program. IZA engages in (i) original and internationally competitive research in all fields of labor economics, (ii) development of policy concepts, and (iii) dissemination of research results and concepts to the interested public.

IZA Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available directly from the author.

ABSTRACT

Do More of Those in Misery Suffer from Poverty, Unemployment or Mental Illness?*

Studies of deprivation usually ignore mental illness. This paper uses household panel data from the USA, Australia, Britain and Germany to broaden the analysis. We ask first how many of those in the lowest levels of life-satisfaction suffer from unemployment, poverty, physical ill health, *and* mental illness. The largest proportion suffer from mental illness. Multiple regression shows that mental illness is not highly correlated with poverty or unemployment, and that it contributes more to explaining the presence of misery than is explained by either poverty or unemployment. This holds both with and without fixed effects.

JEL Classification: I1, I31, I32

Keywords: mental health, life-satisfaction, wellbeing, poverty, unemployment

Corresponding author:

Richard Layard
Centre for Economic Performance
London School of Economics and Political Science
Houghton Street
London WC2A 2AE
United Kingdom
E-mail: R.Layard@lse.ac.uk

* We thank Jörn-Steffen Pischke and George Ward for helpful advice, and Daniel Kahneman, George Lowenstein, Stephen Nickell, Andrew Oswald, Nick Powdthavee and others for their comments. Support from the US National Institute on Aging (Grant R01AG040640) and the UK Economic and Social Research Council is gratefully acknowledged.

Increasingly, policy-makers are considering the life-satisfaction of the population as a possible policy goal.¹ This makes it more important than ever to have a comprehensive account of what determines life-satisfaction. In the current debate most of the factors considered are ‘external’ to the individual – ‘situational’ factors like income, employment, family status, community safety, and religious participation.² The chief ‘internal’ variable that is considered is general health; but this in practice relates mainly to physical health. Mental health is strikingly absent from most empirical analyses of life-satisfaction, and consequently from much of the policy debate. This may help to explain why only 5% of health expenditure in rich countries goes on average to mental health.³

The purpose of this paper is to remedy the omission of mental health from the analysis. The data are household panel data from the USA, Australia, Britain, and Germany. In them, a typical life-satisfaction question is “How satisfied are you with your life, all things considered?”, with possible responses on a numerical scale running from “completely dissatisfied” to “completely satisfied” – what Kahneman calls a measure of evaluated wellbeing.⁴

In our analysis we focus on the lowest levels of life-satisfaction (roughly the bottom 10%) which we call ‘misery’. Directly comparable results on life-satisfaction treated as a continuous variable are given in the online Appendix. Those results are extremely similar to the results on misery, and in themselves represent a significant contribution to the literature on life-satisfaction.

There are many reasons why the relationship between mental health and life-satisfaction has been so widely overlooked in the wellbeing debate. One is the fact that mental illness is a subjective state and so is life-satisfaction. But our data have a key advantage: they include the most “objective” measures available of mental illness – that the person has been diagnosed with depression/anxiety by a health professional, or that the person is in treatment. In addition, because individuals are observed over several years, we can examine how mental illness and misery co-vary within the same lifetime.

Even so, some people might argue that mental illness and misery are the same thing. In this paper we show that, to the contrary, the correlation between misery and mental health (between 0.1 and 0.4) is not high enough to suggest they are measures of the same construct. We find that there are many causes of misery but mental illness is one important cause, even holding constant all other causes.

So treating mental illness directly is one way to reduce misery.⁵ But how large a part should the treatment of mental illness play in a strategy to reduce misery? This depends on how big a problem mental health is compared with other problems: how much of the misery in a society is associated with mental illness, as opposed to issues like poverty, unemployment or physical ill-health? That is what this paper is about.

¹ O'Donnell *et al.* (2014).

² Layard *et al.* (2012).

³ Layard and Clark (2014)

⁴ Kahneman (2011).

⁵ Roth and Fonyagy (2005).

In what follows we begin with the simplest possible, descriptive question: What are the characteristics of the most miserable sections of the community? As we show, many more of those in the lowest levels of life-satisfaction suffer from mental illness than from unemployment, poverty or physical illness. This is true in all four countries. We then move to multivariate analysis where we find that the presence or absence of mental illness explains more of the variance of misery than is explained by either poverty, unemployment or physical illness. This is true in cross-sectional analysis, but also using fixed effects or including the lagged dependent variable.

1. Data and methods

Our analysis uses five household surveys, all of which provide information on both life-satisfaction and on mental health. Three of these surveys have the key advantage of including “objective” data on whether the person has been diagnosed with depression/anxiety, with two of them also including data on whether the person is in treatment for a mental health problem. These three surveys are: for the USA, the Behavioral Risk Factor Surveillance System (BRFSS) and the Panel Study of Income Dynamics (PSID); and for Australia, the Household, Income and Labour Dynamics in Australia (HILDA) Survey.

Of these, the BRFSS is purely cross-sectional, giving data on different people for 2006-2013. But the two other studies have data on the same individual’s diagnostic condition for two different years. This enables us to examine the effect of changes in mental health on changes in life-satisfaction, thus mitigating the influence of unobserved individual characteristics.

However, to obtain multiple repeated observations on the same individual, we have to use data on self-reported symptomatology of mental illness provided by three household panel surveys: for Australia, HILDA annually 2001-2010; for Britain, the British Household Panel Survey (BHPS) annually 1996-2008; and for Germany, the German Socio-Economic Panel (GSOEP) biannually 2002-2008.

All the five surveys provide information on income, employment, family status, education and physical health. Thus we can isolate the impact of mental and physical health on misery, holding constant other potential sources of misery. All the data are from representative samples of people aged 25 and over, and all variables are based on questionnaires set out in full in the Appendix.

1.1. *Mental illness*

Mental illness is measured either by “objective” data (as above) or by self-reported mental health symptomatology (in Australia from the Short Form 36 Health Survey; in the UK from the General Health Questionnaire 12; and in Germany from the Short Form 12 Health Survey). We exclude “happy” items from the Short Form Health Survey and the General Health Questionnaire.

Not all questions are asked in all years and when they are included they are not always answered.⁶ In each analysis we include only observations for which there are replies to all questions, sample sizes being reported in each table.

Physical health is defined in the U.S. by the number of health conditions diagnosed by a health professional; in Australia and Germany by self-reported symptomatology; and in Britain by the self-reported number of conditions. Household income is equivalised. All regression equations also control linearly for age, age², living with a partner, education and gender.

Sample sizes are shown for each regression and means and standard deviations of all variables are shown with the Correlation Matrices in the Appendix.

1.2. *Misery*

We define misery as being in the bottom levels of life-satisfaction. The exact proportion of the population in misery differs between countries because life-satisfaction is measured in discrete integers. Misery in Australia is the bottom 7.5% of life-satisfaction (0-5 on a scale from 0 to 10); in the US (BRFSS) the bottom 5.6% (1-2 on a scale from 1 to 4); in the US (PSID) the bottom 6% (1-2 on a scale from 1 to 5); in Britain the bottom 9.9% (1-3 on a scale from 1 to 7) and in Germany, the bottom 8.7% (0-4 on a scale from 0 to 10).

1.3. *Other variables*


Physical health is defined in the U.S. by the number of health conditions diagnosed by a health professional; in Australia and Germany by self-reported symptomatology; and in Britain by the self-reported number of conditions. Household income is equivalised, with household income per adult equivalent equal to the family income divided by $(1+0.7(\text{other adults})+0.5 \text{ children})$. All regression equations also control linearly for age, age², living with a partner, education and gender – known to be significant predictors of life-satisfaction.

2. Results

2.1. *Descriptive statistics*

We begin in Figure 1 with descriptive statistics, to see what percentage of the people in misery have different specific characteristics.

⁶ Response rates to the questions on diagnosing were BRFSS 88%, PSID 93% and HILDA 64%. For questions on symptomatology the rates were BHPS 92%, GSOEP 95% and HILDA 61%.

Fig. 1: *Percentage of Those in Misery Having the Characteristics Shown*

Notes: Misery in Australia is the bottom 7.5% of life-satisfaction, and in the United States (BRFSS) the bottom 5.6%. In Australia, the sample size is 16,896 (and 8,855 for the question on treatment). In US, the sample size is 217,000-268,000.

In Australia only 20% of the least satisfied segment of the population are poor. The proportion who are unemployed is even smaller. By contrast 48% of those in misery have ever been diagnosed with depression or anxiety disorders, and 31% are currently in treatment. The position in the USA is broadly similar: many more of the least satisfied segment of the population have mental health problems than are poor or unemployed.

What accounts for these differences? Two forces are at work. The first is how likely people with each condition are to be miserable, relative to the likelihood in the general population. Without implying causality, we can call this the “relative impact” of the factor. The second is the overall “prevalence” of the condition in the total population.

Thus, if M_i is the number in misery who have condition i , M is the total number of people in misery, T_i is the total number of people with condition i in the population, and T is the total population, then the numbers in Figure 1 are given by

$$\frac{M_i}{M} \equiv \frac{M_i/T_i}{M/T} \cdot \frac{T_i}{T} \equiv \text{Relative impact} \times \text{Prevalence}$$

Table 1 uses this breakdown to account (arithmetically) for the share of the dissatisfied who have each characteristic.

Table 1
Decomposition of Sources of Misery

	% of those in misery having each characteristic	\equiv	Relative impact of each characteristic on misery	X	% Prevalence of each characteristic
	(1)		(2)		(3)
Australia					
Poor (bottom 10%)	20 (0.4)		2.0 (0.1)		10 (0.1)
Unemployed	7 (0.3)		2.9 (0.1)		2.5 (0.1)
Ever diagnosed with depression/anxiety	48 (1.4)		2.6 (0.1)		18 (0.3)
Currently in treatment for a mental health condition	31 (1.7)		3.4 (0.2)		9 (0.3)
Physical health problems (bottom 10%)	22 (0.5)		2.2 (0.05)		10 (0.1)
United States (BRFSS)					
Poor (bottom 10%)	27 (0.1)		2.7 (0.1)		10 (0.0)
Unemployed	13 (0.1)		3.2 (0.1)		4 (0.0)
Ever diagnosed with depression/anxiety	61 (0.4)		2.7 (0.1)		22 (0.1)
Currently in treatment for a mental health condition	40 (0.4)		3.0 (0.1)		13 (0.1)
Physical health problems (bottom 10%)	14 (0.1)		1.4 (0.0)		10 (0.0)

Notes. Standard errors in parentheses. Misery in Australia is the bottom 7.5% of life-satisfaction and in the United States (BRFSS) the bottom 5.6%. In Australia, the sample size is 16,896 (and 8,855 for the question on treatment). In US, the sample size is 217,000-268,000.

The high share of those who are mentally ill, compared with the share who are poor, is due to a mixture of the greater prevalence of mental illness and its greater relative impact. There is no danger that we have overstated the importance of mental illness since the prevalence of a

mental health diagnosis in these surveys is rather below the prevalence in specific household surveys of psychiatric morbidity.^{7,8,9} The table also suggests that physical health problems (as measured) have a somewhat lower relative impact than mental health problems.

Description is a proper way to start. But an obvious question is whether much of the mental illness is not itself due to poverty and unemployment, in which case the proper policy priority might be to focus on reducing poverty and unemployment rather than directly attacking mental illness. To answer this question requires multivariate analysis.

2.2. Cross-sectional regressions

We choose to use linear multiple regression, with misery as the dependent variable treated as a 0/1 dummy variable. We also use logit analysis which gives almost identical results (shown in the Appendix). But linear multiple regression using standardised variables has a major advantage: the resulting partial correlation coefficients (or β -statistics) shown in Table 2 reflect the “power” of each variable to explain the presence or absence of misery, holding all other variables in the equation constant. They therefore reflect the impact of the variable times its standard deviation, which in the case of a dichotomous variable is $\sqrt{(T_i/T)(1 - T_i/T)}$ where T_i/T is its prevalence.

Table 2
Predictors of Misery: Cross-section Analysis (partial correlation coefficients)

	Australia			United States (BRFSS)		
Income (log)	-0.08 (17)	-0.06 (7)	-0.05 (5)	-0.14 (27)	-0.12 (14)	-0.13 (19)
Unemployed	0.07 (12)	0.06 (5)	0.05 (3)	0.07 (41)	0.06 (18)	0.07 (14)
Ever diagnosed with depression/anxiety		0.14 (14)			0.17 (44)	
Currently in treatment for mental health condition			0.12 (9)			0.16 (39)
Physical health problems	0.17 (34)	0.16 (14)	0.16 (10)	0.09 (35)	0.05 (14)	0.09 (22)
N Observations	81,285	16,896	8,855	1,961,708	268,300	217,225
R ²	0.061	0.090	0.097	0.055	0.084	0.082

Notes. Robust t-statistics in parentheses. Controls for age, age², living with partner, education, and gender.

To increase the explanatory power of income and physical health, we now treat them as continuous variables. Table 2 shows the results. Each column represents one equation. For each country the first equation omits mental health, and therefore shows the standardised ‘effect’ of income and unemployment in a way that includes their effects via mental health. As the t-statistics show, the effects are highly significant. But, when we introduce mental health, the

⁷ McManus *et al.* (2009).

⁸ Kessler *et al.* (2005).

⁹ Wittchen and Jacobi (2005).

effects barely change. This is because of the strikingly low correlation between mental health and income or unemployment. Moreover, once mental health is introduced, it exerts a bigger influence on misery than either income or unemployment.

How about the comparison of mental with physical health? In the table the mental health variables (which are binary) have similar explanatory power to the continuous physical health variables. But it is difficult to draw clear conclusions due to possible problems of measurement. To help with this problem, the BRFSS provided a separate measure of mental health that is exactly analogous to a measure of physical health: each respondent was asked “For how many days during the past 30 days was your mental health not good?”, and an identical question for physical health. The replies to the mental health question had a much stronger predictive power than those on physical health ($\beta = 0.31$ and 0.11 respectively), even though the most miserable people reported almost the same number of days with bad mental and physical health (15 and 13 days respectively). The overall conclusion must be that mental and physical illness have similar power to explain misery, and in both cases greater power than variations in income or employment.

The analysis so far is purely cross-sectional. It barely begins to approach any attempt at causality – for example, the cross-sectional correlations are partly the product of permanent genetic or personality differences affecting both the correlated variables. We can come closer to causality by looking at the same individual at multiple points in time, and examining how changes in different variables within the same person are interconnected.

2.3. Time-series descriptive statistics

We begin with the two surveys which give two years of repeated data on the diagnostic state of the same individual. These are HILDA and the U.S. PSID. (In the PSID the percentage of diagnosed mental illness is only 7%, much smaller than is normal in household surveys of psychiatric morbidity⁷⁻⁹ which is why we have not used it earlier.)

We can start again with simple descriptive statistics. Table 3 examines those people who entered misery within a given period, and asks what else changed in their lives over the same period. In Australia 14% of these people had acquired a diagnosis of depression or anxiety disorder, while only 7% had become poor, 6% had become unemployed, and 8% had become physically ill. In the United States the role of recent poverty in explaining newly-acquired misery was greater than that of mental illness, partly due to the narrow definition of mental illness and partly due to the huge flows in and out of poverty.

Table 3
Decomposition of Sources of Newly-Acquired Misery

	% of those entering misery who have each characteristic	≡	Relative impact of each characteristic upon entering misery	X	% of population who have each characteristic
Australia (2007 to 2009)					
Became poor (bottom 10%)	7.3 (1.3)		1.9 (1.1)		3.7 (0.2)
Became unemployed	5.9 (1.2)		3.5 (1.5)		1.7 (0.1)
Became diagnosed with depression/anxiety	13.5 (2.1)		2.4 (0.7)		5.6 (0.3)
Became physically ill (bottom 10%)	8.4 (1.7)		2.1 (1.0)		3.9 (0.2)
USA (PSID) (2009 to 2011)					
Became poor (bottom 10%)	19.1 (1.5)		3.2 (0.5)		5.9 (0.1)
Became unemployed	11.1 (1.2)		2.3 (0.7)		4.9 (0.1)
Became diagnosed with emotional, nervous or psychiatric problems	13.3 (1.3)		3.9 (0.5)		3.4 (0.1)
Became physically ill (bottom 10%)	8.6 (1.1)		2.7 (0.4)		3.2 (0.1)

Notes. Standard errors in parentheses. Misery in Australia is the bottom 7.5% of life-satisfaction and in the USA (PSID) the bottom 6%. In Australia, the sample size is 16,896. In US, the sample size is 27,095.

2.4. Time-series regressions

However to get nearer to causality we have to look simultaneously at the impact of all variables and to include movements out of misery as well as into it. This is done in Table 4, which is the dynamic equivalent of Table 2. For each country we start with equations including a fixed effect for each individual. This removes the effect of all permanent differences between individuals. When this is done, the coefficients on all variables are substantially reduced, as the comparison for Australia shows. For example, in the case of mental illness we have removed the effect of all permanent differences between individuals in their mental health. But diagnosed mental illness remains a more important explanation of the fluctuation in misery over the life-course than are either income or unemployment. The same is true in the U.S. using the PSID.

Table 4
Predictors of Misery: Dynamic Analysis (partial correlation coefficients)

	Australia		United States (PSID)	
Income (log)	-0.01 (0.3)	-0.03 (4.0)	-0.04 (2.7)	-0.06 (6.0)
Unemployed	0.02 (1.3)	0.05 (3.8)	0.02 (2.3)	0.05 (7.0)
Ever diagnosed with depression/anxiety*	0.04 (1.8)	0.10 (11.5)	0.09 (5.7)	0.08 (10.4)
Physical health problems	0.08 (4.0)	0.11 (12.2)	0.04 (2.6)	0.03 (4.4)
Misery in previous year		0.33 (20.2)		0.24 (29.9)
N Observations	16,896	15,767	27,095	12,450
R ²	0.010	0.186	0.006	0.116
Individual fixed effect	YES	NO	YES	NO

Notes. Robust t-statistics in parentheses. Controls for age, age², living with partner, education, and gender.

* In the U.S. “emotional, nervous or psychiatric problems”.


An alternative way to investigate the dynamics of misery is by replacing the fixed effect by the lagged dependent variable. This takes out less of the fixed effect but also allows for other lagged effects of the observed and unobserved variables. This procedure is used in the remaining columns of Table 4 and again shows mental illness as a more important explanatory factor than income or unemployment.

2.5. Analyses using self-reported symptomatology

The major limitation of the preceding analysis is that we only have data on diagnosed illness for two years. By contrast, if we use self-reported symptoms as the basis for identifying mental illness, we have many more years data from the Australian, British and German household panel surveys (up to 9, 12 and 6 years respectively). Figure 2 shows for these countries the share of misery accounted for by people in the bottom 10% of self-reported mental health. Even if we measure this variable six years earlier, it accounts for more of those in misery than are accounted for by being in poverty or unemployment today.

But again, to attempt to move towards causality, we need to run multiple regressions. Table 5 shows the results of such analysis including fixed effects and treating mental health as a continuous variable. This shows that when we include measures of current symptoms, the estimated effect of mental health is larger than in our previous tables. But this could partly reflect partly time-varying fluctuations in reporting style. When, to obviate this problem, we replace current by lagged mental health, the explanatory power of mental health is less - but most frequently greater than that of current income or employment.

Fig. 2. Percentage of Those in Misery Having the Characteristics Shown



Notes. Those in misery comprise the bottom 7.5% of life-satisfaction in Australia, the bottom 9.9% in Britain, and the bottom 8.7% in Germany.

Table 5
Predictors of Misery: Using Symptomatology and Fixed Effects (partial correlation coefficients)

	Australia		Britain		Germany	
Income (log)	-0.02 (2.8)	-0.01 (1.7)	-0.02 (3.6)	-0.02 (3.5)	-0.03 (5.0)	-0.02 (2.2)
Unemployed	0.03 (5.1)	0.03 (4.1)	0.02 (5.4)	0.03 (6.6)	0.03 (3.6)	0.04 (5.1)
Self-reported mental health problems	0.15 (24.9)		0.33 (60.9)		0.23 (24.8)	
Self-reported mental health problems in previous year	0.02 (3.3)		0.08 (15.0)		0.06 (7.2)	
Physical health problems	0.03 (3.7)	0.06 (7.9)	0.03 (5.6)	0.06 (10.2)	0.04 (5.6)	
Physical health problems in previous year					0.03 (3.5)	
N Observations	81,285	67,003	126,987	113,522	53,407	53,699
N Individuals	15,375	12,652	20,758	18,672	22,673	20,041
R ²	0.029	0.007	0.096	0.009	0.049	0.007
Individual fixed effect	YES	YES	YES	YES	YES	YES

Notes. Robust t-statistics in parentheses. Controls for age, age², living with a partner, education and gender.

3. Discussion

We have shown that, to understand the sources of misery, one should look not only at traditional variables like income, employment and physical health but also at mental health. Though we do not claim anywhere to have a fully causal analysis, our results suggest that mental illness is an important form of deprivation, which receives far too little attention, even within the health sector.¹⁰

Indeed our findings confirm earlier work comparing the effects of physical and mental illness. Here Dolan and Metcalfe regressed life-satisfaction on reported mental and physical pain using the EQ5D, and found that mental pain had a bigger effect than physical pain.¹¹ That study was cross-sectional (as was another¹²), but in a further study Dolan *et al.* repeated the analysis using two separate observations per person, with similar results.¹³ The authors hypothesised that mental pain is more difficult to adapt to – it occupies more of a person’s mental space. Like their work, our results strongly suggest that health policy should give greater weight to mental health and that the weights used in calculating QALYs should be reconsidered.¹⁴

However when it comes to policy-making, two more key questions are the efficacy of treatment and the cost. To compare these, the policy-maker will want to evaluate their effects on life-satisfaction, which is a preferred policy measure to ‘misery’.¹ So how does treatment increase life-satisfaction, and how does cost reduce it?

We can illustrate this by considering the case for better access to cognitive behavioural therapy for people with depression or anxiety disorders. We shall use coefficients from the life-satisfaction regressions shown in the Appendix. On the benefit side these show that, holding constant the fixed effect, acquiring a diagnosis in the USA reduces life-satisfaction by 0.3 standard deviations ($-0.07/-0.26$). So losing a diagnosis raises life-satisfaction by an equal amount, and from field trials we know that therapy leads at least 1 in 3 to recover who would not otherwise have done so.⁵ So treatment yields an average gain of 0.1 standard deviations of life-satisfaction per person treated, lasting for at least a year.

This has to be compared with the cost. The therapy costs some 5% of average annual income per person, which is about 0.05 standard deviations of annual equivalised log income in the USA. This reduces annual life-satisfaction by 0.0015 standard deviations (0.05×0.03). This compares with the benefit of 0.1 standard deviations: the benefit is some 70 times the cost.

In practice the cost would of course be spread across more people than those who benefit from the treatment, but this spreading of the cost would only serve to reduce its total impact. For Australia the ratio of benefit to cost is about 10 times. These calculations are extremely rough but simply illustrate how this approach could provide policy-makers with a quite new perspective on orders of magnitude.

¹⁰ Another strand of research has examined the relation between life-satisfaction and personality factors, including neuroticism and extroversion, see Diener and Seligman (2002); Graham *et al.* (2009); Vazquez *et al.* (2014); Headey *et al.* (1993); Boyce *et al.* (2013).

¹¹ Dolan and Metcalfe (2012)

¹² Mukuria and Brazier (2013)

¹³ Dolan *et al.* (2012)

¹⁴ Dolan and Kahneman (2008).

The analysis of the paper is of course subject to a host of caveats. The analysis is not fully causal, though it becomes more so through the inclusion of fixed effects or the lagged dependent variable. Moreover, question-ordering can introduce spurious correlation between variables.¹⁵ For example, in the BHPS mental health questions precede the question on life-satisfaction and the former may influence the latter. But the BRFSS, PSID and GSOEP ask about life-satisfaction before mental health, and HILDA asks the question on separate occasions. Moreover in fixed effects analysis the bias resulting from question-ordering largely disappears provided the question-ordering does not change.

We conclude that time-varying mental health really has an important independent influence on life-satisfaction. And so surely does the non-varying component of mental health, though this is less easily studied. There are policy implications. For too long the debate on deprivation has focussed mainly on poverty, jobs, education and physical sickness. It needs broadening to include the inner person. This is the new frontier for labour economics.¹⁶

Additional Supporting Information may be found in the online version of this article:

Appendix A. Data sources.

Appendix B. Definitions of variables.

Appendix C. Replication of text table 1 for PSID.

Table C1. Decomposition of sources of misery

Appendix D. Replication of text tables 2, 4 and 5 with life-satisfaction as the dependent variable.

Table D1. Predictors of life-satisfaction: Cross-section analysis

Table D2. Predictors of life-satisfaction: Dynamic analysis

Table D3. Predictors of life-satisfaction: Using symptomatology and fixed effects.

Appendix E. Correlation matrices, mean and standard deviations.

Appendix F. Logit analysis.

¹⁵ Schwarz and Strack (1999).

¹⁶ Layard (2014).

References

- Boyce, C.J., Wood, A.M. and Powdthavee, N. (2013). 'Is personality fixed? Personality changes as much as "variable" economic factors and more strongly predicts changes to life satisfaction', *Social Indicators Research*, vol. 111(1), pp. 287-305.
- Diener, E. and Seligman, M. (2002). 'Very happy people', *Psychological Science*, vol. 13(1), pp. 81-84.
- Dolan, P. and Kahneman, D. (2008). 'Interpretations of utility and their implications for the valuation of health', *Economic Journal*, vol. 118(525), pp. 215-234.
- Dolan P, Lee H. and Peasgood T. (2012). 'Losing sight of the wood for the trees: Some issues in describing and valuing health, and another possible approach', *Pharmacoeconomics*, vol. 30(11), pp. 1035-49.
- Dolan P. and Metcalfe R. (2012). 'Valuing health: a brief report on subjective well-being versus preferences', *Medical Decision Making*, vol. 32(4), pp. 578-82.
- Graham, C., Jaramillo, L.F.H. and Lora, E.A. (2009). *Valuing health conditions: Insights from happiness surveys across countries and cultures. Working Paper 4625*. Washington, D.C, Inter-American Development Bank.
- Headey, B., Kelley, J. and Wearing, A. (1993). 'Dimensions of mental health: Life satisfaction, positive affect, anxiety and depression', *Social Indicators Research*, vol. 29, pp. 63-82.
- Kahneman, D. (2011). *Thinking, fast and slow*, London: Allen Lane.
- Kessler, R.C., Chiu, W.T., Demler, O., Merikangas, K.R. and Walters, E.E. (2005). 'Prevalence, severity, and comorbidity of 12-month *DSM-IV* disorders in the National Comorbidity Survey Replication', *Archives of General Psychiatry*, vol. 62(6), pp. 617-627.
- Layard, R. (2014). 'Mental Health: the new frontier for labour economics', in (D. McDaid and C.L. Cooper, eds.), *Wellbeing: A complete reference guide Volume 5 (The economics of wellbeing)*, pp.: Wiley-Blackwell.
- Layard R. and Clark DM. (2014). *Thrive: the power of evidence-based psychological therapies*. London: Penguin.
- Layard, R., Clark, A.E. and Senik, C. (2012). 'The causes of happiness and misery', in (J.F. Helliwell, R. Layard and J. Sachs, eds.), *World Happiness Report*, pp. 58-89, New York: The Earth Institute, Columbia University.
- McManus, S., Meltzer, H., Brugha, T., Bebbington, P. and Jenkins, R., Eds. (2009). *Adult psychiatric morbidity in England, 2007: results of a household survey*. Leeds: The Health & Social Care Information Centre, Social Care Statistics.
- Mukuria C., and Brazier J. (2013). 'Valuing the EQ-5D and the SF-6D health states using subjective well-being: A secondary analysis of patient data', *Social Science & Medicine*, vol. 77, pp. 97-105.
- O'Donnell, G., Deaton, A., Durand, M., Halpern, D. and Layard, R. (2014). *Wellbeing and policy*. London: Legatum Institute.
- Roth, A. and Fonagy, P., Eds. (2005). *What works for whom? A critical review of psychotherapy research*. Second edition. New York: Guilford Press.
- Schwarz, N. and Strack, F. (1999). 'Reports of Subjective Well-Being: Judgmental Processes and Their Methodological Implications', in (D. Kahneman, E. Diener and N. Schwarz, eds.), *Well-Being: The Foundations of Hedonic Psychology*, pp., New York: Russell Sage Foundation.
- Vazquez, C., Rahona, J.J., Gomez, D., Caballero, F.F. and Hervas, G. (2014). 'A National Representative Study of the Relative Impact of Physical and Psychological Problems on Life Satisfaction', *Journal of Happiness Studies*, Published online 1 February.

Wittchen, H.-U. and Jacobi, F. (2005). 'Size and burden of mental disorders in Europe: a critical review and appraisal of 27 studies', *European Neuropsychopharmacology*, vol. 15, pp. 357-376.

Appendix

- A. Data sources**
- B. Definitions of variables**
- C. Replication of text table 1 for PSID**
- D. Replication of text tables 2, 4 and 5 with life-satisfaction as the dependent variable**
- E. Correlation matrices, mean and standard deviations**
- F. Logit analysis**

A. Data sources

<p>Australia</p>	<p>Household Income and Labour Dynamics in Australia (HILDA) Survey</p> <p>Household-based panel study which began in 2001. The panel members are followed over time and interviewed every year. Life-satisfaction is measured throughout.</p> <p>In 2007 and 2009, respondents were asked whether they have ever been diagnosed with depression or anxiety.</p> <p>In 2009, they were also asked whether they take prescription medication for depression or anxiety or whether they have been seen during the last 12 months by a mental health professional.</p> <p>From 2001 to 2010, the SF-36 questionnaire is included.</p>
<p>USA (BRFSS)</p>	<p>Behavioral Risk Factor Surveillance System (BRFSS)</p> <p>Cross-sectional survey which includes a life-satisfaction question since 2005.</p> <p>There are several measures of mental health in the BRFSS: Ever diagnosed with depression or anxiety: 2006; 2008; 2010; 2013 Receiving mental health treatment: 2007; 2009 Days mental health not good this month: 2005-2010; 2013</p>
<p>USA (PSID)</p>	<p>Panel Study of Income Dynamics (PSID)</p> <p>Household-based panel study. A life-satisfaction question is included in 2009 and 2011.</p> <p>In 2009 and 2011, respondents were also asked whether they have ever been diagnosed with depression or anxiety.</p>
<p>Britain</p>	<p>British Household Panel Survey (BHPS)</p> <p>Household-based panel study which began in 1991. The panel members are followed over time and interviewed every year. A life satisfaction question has been included in the study from 1996.</p> <p>From 1996 to 2008, it also collects information on mental health using the GHQ-12 questionnaire.</p>
<p>Germany</p>	<p>German Socio Economic Panel (GSOEP)</p> <p>Household-based panel study which began in 1984. The panel members are followed over time and interviewed every year. Mental and physical health are measured using the SF-12 questionnaire in 2002; 2004; 2006 and 2008. Life-satisfaction is measured throughout.</p>

B. Definitions of variables

Life Satisfaction	
Australia	All things considered, how satisfied are you with your life? Pick a number between 0 and 10 to indicate how satisfied you are.
USA (BRFSS)	In general, how satisfied are you with your life? 1-4
USA (PSID)	Please think about your life-as-a whole. How satisfied are you with it? 1-5
Britain	How dissatisfied or satisfied are you with your life overall? 1-7
Germany	How satisfied are you with your life, all things considered? 0-10
Mental health measures	
(1) Diagnosis	
Australia	Have you ever been told by a doctor or nurse that you have any long term health conditions listed below? eg. Depression/Anxiety. (Yes/No)
USA (BRFSS)	Yes to either or both of the following: <ul style="list-style-type: none"> • Have you ever been told you have an anxiety disorder? (Yes/No) • And/or have you ever been told you had a depressive disorder? (Yes/No)
USA (PSID)	Has a doctor or other health professional ever told you that you had any emotional, nervous, or psychiatric problems? (Yes/No)
(2) Treatment	
Australia	Yes to either or both of the following: <ul style="list-style-type: none"> • Takes prescription medication for depression or anxiety. (Yes/No) • Seen during last 12 months a mental health professional (Yes/No)
USA (BRFSS)	Are you now taking medicine or receiving treatment from a doctor or other health professional for any type of mental health condition or emotional problem? (Yes/No)
(3) Days mental health not good this month	
USA (BRFSS)	Now thinking about your mental health, which includes stress, depression and problems with emotions, for how many days during the past 30 days was your mental health not good? (0/30)
(4) Self-reported symptomatology	
Australia	<p>SF-36 questionnaire</p> <p>Mental health: These questions are about how you feel and how things have been with you during the past 4 weeks. For each question, please file the one answer that comes closest to the way you have been feeling. How much of the time during the past 4 weeks:</p> <ul style="list-style-type: none"> • Have you been a nervous person? (1-5) • Have you felt so down in the dumps that nothing could cheer you up? (1-5) • Have you felt calm and peaceful? (1-5) • Have you felt down? (1-5) • Have you been a happy person? (1-5) <p>Emotional factors affecting role: During the past 4 weeks, have you had any of the following problems with your work or other regular daily activities as a result of any emotional problems (such as feeling depressed or anxious)?</p> <ul style="list-style-type: none"> • Cut down the amount of time you spent on work or other activities (1-3) • Accomplished less than you would like (1-3) • Did not do work or other activities as carefully as usual (1-3) <p>To calculate our self-reported mental health measure, we first create the “mental health” and the “emotional factors affecting role” indicators that are total score from the above questions, each of them rescaled from 1 to</p>

	<p>5. Then, we compute the total score of these two new variables to obtain the self-reported mental health measure.</p>
<p>Britain</p>	<p>General Health Questionnaire-12 Number of Yes answers.</p> <p>Here are some questions regarding the way you have been feeling over the last few weeks. For each question please tick the box next to the answer that best describes the way you have felt.</p> <ul style="list-style-type: none"> • Have you recently been able to concentrate on whatever you are doing? (Yes/No) • Have you recently lost much sleep over worry? (Yes/No) • Have you recently felt that you were playing a useful part in things? (Yes/No) • Have you recently felt capable of making decisions about things? (Yes/No) • Have you recently felt constantly under strain? (Yes/No) • Have you recently felt you could not overcome your difficulties? (Yes/No) • Have you recently been able to enjoy your normal day-to-day activities? (Yes/No) • Have you recently been able to face up to problems? (Yes/No) • Have you recently been feeling unhappy or depressed? (Yes/No) • Have you recently been losing confidence in yourself? (Yes/No) • Have you recently been thinking of yourself as a worthless person? (Yes/No) • Have you recently been feeling reasonably happy, all things considered? (Yes/No)
<p>Germany</p>	<p>SF-12 questionnaire</p> <p>Please think about the last four weeks. How often did it occur within this period of time:</p> <p>Mental health:</p> <ul style="list-style-type: none"> • That you felt run-down and melancholy? (1-5) • That you felt relaxed and well-balanced? (1-5) <p>Emotional factors affecting role:</p> <ul style="list-style-type: none"> • That due to mental health or emotional health problems: <ul style="list-style-type: none"> ○ You achieved less than you wanted to at work or in everyday tasks? (1-3) ○ You carried out your work or everyday tasks less thoroughly than usual? (1-3) <p>To calculate our self-reported mental health measure, we first create the “mental health” and the “emotional factors affecting role” indicators that are total score from the above questions, each of them rescaled from 1 to 5. Then, we compute the total score of these two new variables to obtain the self-reported mental health measure.</p>

Physical health measures	
(1) Number of physical health problems	
USA (BRFSS)	<p>Number of Yes answers.</p> <p>Have you ever been told by a doctor, nurse or other health professional, that you had:</p> <ul style="list-style-type: none"> • Diabetes (Yes/No) • Heart attack (Yes/No) • Angina or coronary heart disease (Yes/No) • Stroke (Yes/No) • Asthma (Yes/No) • Arthritis (Yes/No) • Cataracts (Yes/No) • Glaucoma (Yes/No) • Macular degeneration (Yes/No) • Prostate cancer (Yes/No)
USA (PSID)	<p>Number of Yes answers.</p> <p>Has a doctor ever told you that you have or had any of the following:</p> <ul style="list-style-type: none"> • Stroke (Yes/No) • Heart attack (Yes/No) • Heart disease (Yes/No) • Hypertension (Yes/No) • Asthma (Yes/No) • Lung disease (Yes/No) • Diabetes (Yes/No) • Arthritis (Yes/No) • Memory Loss (Yes/No) • Learning Disorder (Yes/No) • Cancer (Yes/No)
Britain	<p>Number of Yes answers.</p> <p>Do you have any of the health problems or disabilities listed on this card? Exclude temporary conditions.</p> <ul style="list-style-type: none"> • Problems or disability connected with: arms, legs, hands, feet, back, or neck (including arthritis and rheumatism) (Yes/No) • Difficult in seeing (other than needing glasses to read normal size print) (Yes/No) • Difficulty in hearing (Yes/No) • Skin conditions/allergies (Yes/No) • Chest/breathing problems, asthma, bronchitis (Yes/No) • Heart/blood pressure or blood circulation problems (Yes/No) • Stomach/liver/kidneys or digestive problems (Yes/No) • Diabetes (Yes/No) • Epilepsy (Yes/No) • Migraine or frequent headaches (Yes/No) • Cancer (Yes/No) • Stroke (Yes/No)

(2) Days physical health not good this month	
USA (BRFSS)	Now thinking about your physical health, which includes physical illness and injury, for how many days during the past 30 days was your physical health not good? (0/30)
(3) Self-reported symptomatology	
Australia	<p>SF-36 questionnaire</p> <p>Physical functioning The following questions are about activities you might do during a typical day. Does your health now limit you in these activities? If so, how much?</p> <ul style="list-style-type: none"> • Vigorous activities, such as running, lifting heavy objects, participating in strenuous sports (1-3) • Moderate activities, such as moving a table, pushing a vacuum cleaner, bowling or playing golf (1-3) • Lifting or carrying groceries (1-3) • Climbing several flights of stairs (1-3) • Climbing one flight of stairs (1-3) • Bending, kneeling, or stooping (1-3) • Walking more than one kilometre (1-3) • Walking half a kilometre (1-3) • Walking 100 metres (1-3) • Bathing or dressing yourself (1-3) <p>Physical factors affecting role During the past 4 weeks, have you had any of the following problems with your work or other regular daily activities as a result of your physical health?</p> <ul style="list-style-type: none"> • Cut down the amount of time you spent on work or other activities (1-5) • Accomplished less than you would like (1-5) • Were limited in the kind of work or other activities (1-5) • Had difficulties performing the work or other activities (1-5) <p>Bodily pain</p> <ul style="list-style-type: none"> • How much bodily pain have you had during the past 4 weeks? (1-5) • During the past 4 weeks, how much did pain interfere with your normal work (including both work outside the home and housework)? (1-5) <p>To calculate our self-reported physical health measure, we first create the “physical functioning”, the “bodily pain” and the “physical factors affecting role” indicators that are total score from the above questions, each of them rescaled from 1 to 5. Then, we compute the total score of these three new variables to obtain the self-reported physical health measure.</p>
Germany	<p>SF-12 questionnaire</p> <p>Physical functioning</p> <ul style="list-style-type: none"> • When you ascend stairs, i.e. go up several floors on foot: Does your state of health affect you greatly, slightly or not at all? (1-3) • And what about having to cope with other tiring everyday tasks, i.e. when one has to lift something heavy or when one requires agility: Does your state of health affect you greatly, slightly or not at all ? (1-3)

	<p>Physical factors affecting role Please think about the last four weeks. How often did it occur within this period of time,</p> <ul style="list-style-type: none"> • That due to physical health problems <ul style="list-style-type: none"> ○ You achieved less than you wanted to at work or in everyday tasks? (1-5) ○ You were limited in some form at work or in everyday tasks? (1-5) <p>Bodily pain</p> <ul style="list-style-type: none"> • That you had strong physical pains? (1-5) <p>To calculate our self-reported physical health measure, we first create the “physical functioning”, the “bodily pain” and the “physical factors affecting role” indicators that are total score from the above questions, each of them rescaled from 1 to 5. Then, we compute the total score of these three new variables to obtain the self-reported physical health measure.</p>
Income (equivalised; OECD scale)	
Australia	Household financial year disposable income
USA (BRFSS)	Gross annual household income from all sources (0-8). All ranges are valued at mid-point, except for top range valued at 1.5 times its lowest value.
USA (PSID)	Total family income (before tax): this variable includes taxable, transfer and social security incomes.
Britain	Gross annual household income. This is the sum of labour and non-labour income.
Germany	Pre-government annual household income: this variable is the sum of total family income from labour earnings, asset flows, private retirement income and private transfers. Labour earnings include wages and salary from all employment, including training, self-employment income and bonuses, overtime and profit-sharing. Asset flows include income from interest, dividends and rent. Private transfers include payments from individuals outside of the household including alimony and child support payments.
Age	<p>We restrict the sample for people aged 25 and over.</p> <p>Age is measured by an aggregate of age and age squared from a previous regression.</p>
Education	Education is measured by an index (with weights from a previous regression).
Having a partner	Having a partner is equal to 1 if respondent is married or cohabiting. It is 0 otherwise.

C. Replication of text table 1 for PSID

Table C1
Decomposition of Sources of Misery

	% of those in misery having each characteristics	≡	Relative impact of each characteristic upon misery	x	% of population who have each characteristic
	(1)		(2)		(3)
United States (PSID)					
In poverty (bottom 10%)	22 (0.8)		2.2 (0.1)		10 (0.1)
Unemployed	14 (0.7)		2.1 (0.1)		7 (0.1)
Ever diagnosed with emotional, nervous or psychiatric problems	20 (0.8)		2.8 (0.1)		7 (0.1)
Physical health problems (bottom 10%)	16 (0.7)		1.6 (0.1)		10 (0.1)

Notes: Standard errors are in parentheses. In this table, as in the main text table, the numbers in each cell in columns (1) and (3) are based on those for whom we have responses on the characteristics in question. For example the figure 22% at the top of column (1) relates to those in misery for whom we have their income. Similarly the figure 10% at the top of column (3) relates to everyone for whom we have their income. Column (2) is obtained as column (1) divided by column (3). Its standard error is taken from a regression equation.

D. Replication of text tables 2, 4 and 5 with life satisfaction as the dependent variable

Table D1
Predictors of Life-Satisfaction: Cross-section Analysis (partial correlation coefficients)

	Australia			United States (BRFSS)		
Income (log)	0.05 (13)	0.04 (5)	0.04 (3)	0.18 (36)	0.16 (23)	0.17 (26)
Unemployed	-0.05 (11)	-0.05 (5)	-0.04 (3)	-0.06 (44)	-0.06 (17)	-0.06 (15)
Ever diagnosed with depression or anxiety		-0.16 (20)			-0.20 (60)	
Currently in treatment for mental health condition			-0.13 (12)			-0.17 (62)
Physical health problems	-0.21 (44)	-0.19 (19)	-0.20 (14)	-0.11 (46)	-0.06 (15)	-0.11 (32)
Age	0.30 (63)	0.26 (26)	0.26 (19)	0.10 (33)	0.06 (17)	0.10 (24)
Married	0.15 (42)	0.14 (19)	0.15 (15)	0.19 (70)	0.16 (45)	0.18 (40)
Educated	0.04 (11)	0.03 (4)	0.02 (2)	0.05 (29)	0.06 (15)	0.06 (21)
Female	0.11 (17)	0.13 (10)	0.12 (7)	0.04 (13)	0.09 (14)	0.07 (11)
N Observations	81,285	16,896	8,855	1,961,708	268,300	217,225
R ²	0.103	0.138	0.139	0.108	0.146	0.136

Notes: Robust t-statistics in parentheses.

Table D2
Predictors of Life-Satisfaction: Dynamic Analysis (partial correlation coefficients)

	Australia			United States (PSID)	
Income (log)	0.04 (2.2)	0.02 (2.7)	0.01 (0.9)	0.03 (2.0)	0.04 (4.4)
Unemployed	-0.03 (2.3)	-0.03 (4.2)	-0.03 (2.1)	-0.03 (4.2)	-0.05 (6.8)
Ever diagnosed with depression or anxiety*	-0.03 (1.8)	-0.09 (18)		-0.07 (4.8)	-0.08 (9.9)
Currently in treatment for mental health condition			-0.09 (8.6)		
Physical health problems	-0.09 (5.4)	-0.10 (12)	-0.10 (8.6)	-0.06 (4.4)	-0.07 (8.8)
Age	0.00 (0.0)	0.12 (12)	0.11 (9.2)	0.65 (9.4)	0.04 (5.2)
Married	0.12 (4.1)	0.07 (10)	0.07 (7.5)	0.11 (6.5)	0.12 (14)
Educated		0.02 (3.3)	0.02 (1.7)		0.01 (0.9)
Female		0.07 (5.6)	0.06 (3.5)		0.03 (2.1)
Life Satisfaction in previous year		0.52 (53)	0.53 (41)		0.39 (48)
N Observations	16,896	15,767	8,178	27,095	12,450
R ²	0.015	0.376	0.380	0.015	0.237
Individual fixed effect	YES	NO	NO	YES	NO

Notes: Robust t-statistics in parentheses.

*In the US, “emotional, nervous or psychiatric problems”.

Table D3
Predictors of Life-satisfaction: Using Symptomatology and Fixed Effects (partial correlation coefficients)

	Australia		Britain		Germany	
Income (log)	0.02 (4.3)	0.02 (3.3)	0.01 (3.1)	0.01 (3.3)	0.04 (8.0)	0.03 (3.4)
Unemployed	-0.02 (5.2)	-0.02 (4.8)	-0.01 (4.2)	-0.02 (6.4)	-0.02 (4.4)	-0.03 (6.2)
Self-reported mental health problems	-0.18 (38)		-0.37 (99)		-0.28 (41)	
Self-reported mental health problems in previous year	-0.03 (6.5)		-0.08 (21)		-0.09 (14)	
Physical health problems	-0.04 (6.2)	-0.08 (12)	-0.03 (8.3)	-0.07 (14)	-0.08 (12)	
Physical health problems in previous year					-0.01 (2.3)	
Age	0.17 (9.7)	0.16 (7.2)	0.10 (7.8)	0.12 (7.5)	0.25 (11)	0.33 (14)
Married	0.12 (15)	0.13 (13)	0.07 (12)	0.08 (11)	0.04 (4.8)	0.04 (4.0)
N Observations	81,285	67,003	126,987	113,522	53,407	53,699
N Individuals	15,375	12,652	20,758	18,672	22,673	20,041
R ²	0.060	0.017	0.170	0.018	0.115	0.019
Individual fixed effect	YES	YES	YES	YES	YES	YES

Notes: Robust t-statistics in parentheses.

E. Correlation Matrices, Means and Standard Deviations

Table E1: *Australia*

	1	2	3	4	5	6	7	8	9	10	11	12
1.Life Satisfaction	1	-	-0.24	-0.22	-0.15	-0.11	0.00	-0.08	0.14	0.15	0.04	0.06
2. Misery	-	1	0.21	0.21	0.18	0.15	-0.10	0.09	-0.01	0.13	0.00	0.04
3.Ever Diagnosed with depression or anxiety	-0.24	0.21	1	0.57	0.29	0.18	-0.09	0.06	-0.02	-0.11	0.10	0.03
4.Currently in treatment for mental health condition	-0.22	0.21	0.57	1	0.24	0.17	-0.08	0.05	-0.01	-0.09	0.07	0.03
5.Self-reported mental health	-0.15	0.18	0.29	0.24	1	0.24	-0.15	0.07	0.03	-0.12	0.05	0.10
6.Self-reported physical health	-0.11	0.15	0.18	0.17	0.24	1	-0.35	0.01	0.44	-0.13	0.06	0.21
7.Income (log)	0.00	-0.10	-0.09	-0.08	-0.15	-0.35	1	-0.05	-0.47	0.15	-0.08	-0.33
8.Unemployed	-0.08	0.09	0.06	0.05	0.07	0.01	-0.05	1	-0.08	-0.07	-0.03	0.01
9.Age	0.14	-0.01	-0.02	-0.01	0.03	0.44	-0.47	-0.08	1	-0.08	0.02	0.29
10.Married	0.15	-0.13	-0.11	-0.09	-0.12	-0.13	0.15	-0.07	-0.08	1	-0.08	-0.07
11.Female	0.04	0.00	0.10	0.07	0.05	0.06	-0.08	-0.03	0.02	-0.08	1	0.13
12.Educated	0.06	0.04	0.03	0.03	0.10	0.21	-0.33	0.01	0.29	-0.07	0.13	1
Mean	7.88	0.08	0.18	0.09	17.65	23.00	9.91	0.02	49.38	0.71	0.53	0.34
SD	1.53	0.26	0.38	0.29	2.56	5.00	0.77	0.16	15.76	0.45	0.50	0.47

Table E2: *USA (BRFSS)*

	1	2	3	4	5	6	7	8	9	10	11	12	13
1. Life Satisfaction	1.00	-0.62	-0.10	-0.36	-0.24	-0.25	-0.21	0.23	-0.11	0.03	0.21	-0.01	0.14
2. Misery	-0.62	1.00	0.09	0.36	0.23	0.22	0.19	-0.16	0.10	-0.03	-0.13	0.01	-0.07
3. Physical health problems	-0.10	0.09	1.00	0.11	0.19	0.12	0.13	-0.03	0.01	0.16	-0.07	-0.11	-0.06
4. Days mental health not good	-0.36	0.36	0.11	1.00	0.33	0.35	0.34	-0.19	0.09	-0.09	-0.10	0.07	-0.09
5. Days physical health not good	-0.24	0.23	0.19	0.33	1.00	0.21	0.20	-0.22	0.03	0.12	-0.11	0.04	-0.13
6. Ever diagnosed with depression or anxiety	-0.25	0.22	0.12	0.35	0.21	1.00	--	-0.11	0.06	-0.07	-0.09	0.12	-0.03
7. Currently in treatment for mental health condition	-0.21	0.19	0.13	0.34	0.20	--	1.00	-0.09	0.03	-0.02	-0.08	0.09	-0.03
8. Income (log)	0.23	-0.16	-0.03	-0.19	-0.22	-0.11	-0.09	1.00	-0.17	0.02	0.17	-0.10	0.38
9. Unemployed	-0.11	0.10	0.01	0.09	0.03	0.06	0.03	-0.17	1.00	-0.10	-0.06	-0.01	-0.05
10. Age	0.03	-0.03	0.16	-0.09	0.12	-0.07	-0.02	0.02	-0.10	1.00	-0.16	0.02	-0.12
11. Married	0.21	-0.13	-0.07	-0.10	-0.11	-0.09	-0.08	0.17	-0.06	-0.16	1.00	-0.12	0.11
12. Female	-0.01	0.01	-0.11	0.07	0.04	0.12	0.09	-0.10	-0.01	0.02	-0.12	1.00	-0.06
13. Educated	0.14	-0.07	-0.06	-0.09	-0.13	-0.03	-0.03	0.38	-0.05	-0.12	0.11	-0.06	1.00
Mean	3.39	0.06	0.17	3.31	4.35	0.22	0.13	10.0	0.04	55.7	0.57	0.62	0.34
SD	0.63	0.23	0.13	7.63	8.87	0.41	0.34	0.80	0.20	15.8	0.49	0.48	0.47

Table E3: USA (PSID)

	1	2	3	4	5	6	7	8	9	10
1. Life Satisfaction	1.00	-0.55	-0.14	-0.13	0.15	-0.08	0.03	0.22	-0.00	0.09
2. Misery	-0.55	1.00	0.12	0.10	-0.12	0.07	0.00	-0.12	-0.00	-0.08
3. Ever diagnosed with emotional, nervous or psychiatric problems	-0.14	0.12	1.00	0.21	-0.12	0.02	-0.02	-0.10	0.01	-0.02
4. Physical health problems	-0.13	0.10	0.21	1.00	-0.07	0.01	0.29	-0.05	-0.01	-0.10
5. Income (log)	0.15	-0.12	-0.12	-0.07	1.00	-0.15	0.04	0.37	-0.03	0.39
6. Unemployed	-0.08	0.07	0.02	0.01	-0.15	1.00	-0.13	-0.11	-0.02	-0.11
7. Age	0.03	0.00	-0.02	0.29	0.04	-0.13	1.00	0.09	0.02	-0.12
8. Married	0.22	-0.12	-0.10	-0.05	0.37	-0.11	0.09	1.00	-0.06	0.14
9. Female	-0.00	-0.00	0.01	-0.01	-0.03	-0.02	0.02	-0.06	1.00	0.04
10. Educated	0.09	-0.08	-0.02	-0.10	0.39	-0.11	-0.12	0.14	0.04	1.00
Mean	3.80	0.06	0.07	1.69	9.91	0.07	45.83	0.56	0.52	13.18
SD	0.87	0.22	0.26	1.28	0.98	0.25	15.16	0.50	0.50	2.44

Table E4: Britain

	1	2	3	4	5	6	7	8	9	10
1. Life Satisfaction	1.00	-0.70	-0.56	-0.16	0.05	-0.08	0.11	0.13	-0.00	-0.01
2. Misery	-0.70	1.00	0.45	0.16	-0.09	0.08	-0.01	-0.11	0.02	-0.06
3. Self-reported mental health	-0.56	0.45	1.00	0.23	-0.08	0.05	0.01	-0.07	0.12	-0.08
4. Physical health problems	-0.16	0.16	0.23	1.00	-0.15	-0.01	0.39	-0.12	0.08	-0.23
5. Income (log)	0.05	-0.09	-0.08	-0.15	1.00	-0.13	-0.18	0.08	-0.06	0.34
6. Unemployed	-0.08	0.08	0.05	-0.01	-0.13	1.00	-0.08	-0.06	-0.05	-0.04
7. Age	0.11	-0.01	0.01	0.39	-0.18	-0.08	1.00	-0.15	0.02	-0.38
8. Married	0.13	-0.11	-0.07	-0.12	0.08	-0.06	-0.15	1.00	-0.10	0.11
9. Female	-0.00	0.02	0.12	0.08	-0.06	-0.05	0.02	-0.10	1.00	-0.07
10. Educated	-0.01	-0.06	-0.08	-0.23	0.34	-0.04	-0.38	0.11	-0.07	1.00
Mean	5.23	0.10	23.4	1.20	9.5	0.03	49.93	0.72	0.55	0.28
SD	1.30	0.30	5.47	1.31	0.75	0.16	16.3	0.45	0.50	0.45

Table E5: *Germany*

	1	2	3	4	5	6	7	8	9	10
1. Life Satisfaction	1.00	-0.67	-0.51	-0.36	0.11	-0.10	-0.03	0.08	-0.00	0.06
2. Misery	-0.67	1.00	0.37	0.24	-0.07	0.08	0.03	-0.06	0.00	-0.03
3. Self-reported mental health	-0.51	0.37	1.00	0.48	-0.08	0.04	0.02	-0.07	0.12	-0.07
4. Self-reported physical health	-0.36	0.24	0.48	1.00	-0.32	0.01	0.43	0.01	0.09	-0.19
5. Income (log)	0.11	-0.07	-0.08	-0.32	1.00	-0.02	-0.49	0.06	-0.08	0.22
6. Unemployed	-0.10	0.08	0.04	0.01	-0.02	1.00	-0.08	-0.02	-0.02	-0.03
7. Age	-0.03	0.03	0.02	0.43	-0.49	-0.08	1.00	0.03	0.03	-0.14
8. Married	0.08	-0.06	-0.07	0.01	0.06	-0.02	0.03	1.00	-0.07	-0.03
9. Female	-0.00	0.00	0.12	0.09	-0.08	-0.02	0.03	-0.07	1.00	-0.05
10. Educated	0.06	-0.03	-0.07	-0.19	0.22	-0.03	-0.14	-0.03	-0.05	1.00
Mean	6.95	0.09	8.54	11.64	8.94	0.05	49.16	0.70	0.52	0.32
SD	1.84	0.28	3.03	4.50	0.87	0.22	15.47	0.46	0.50	0.47

F. Logit analysis

Table F6
Predictor of Misery: Logit Analysis (partial correlation coefficients)

	Australia			United States (BRFSS)				
	Misery	Misery	Misery	Misery	Misery	Misery	Misery	Misery
Income (log)	-0.272 (18.3)	-0.206 (6.2)	-0.191 (4.5)	-0.494 (35.2)	-0.438 (17.3)	-0.428 (22.3)	-0.289 (25.7)	-0.413 (16.0)
Unemployed	0.141 (13.9)	0.133 (5.4)	0.114 (3.4)	0.146 (41.1)	0.145 (21.2)	0.157 (17.3)	0.145 (42.5)	0.152 (21.5)
Ever diagnosed with depression or anxiety		0.447 (16.2)			0.624 (67.3)			
Currently in treatment for mental health conditions			0.305 (10.8)			0.446 (46.4)		
Days mental health not good							0.649 (87.5)	
Days physical health not good							0.307 (83.8)	
Ever diagnosed with anxiety disorder								0.219 (25.7)
Ever diagnosed with depressive disorder								0.518 (51.9)
Physical health problems	0.556 (41.8)	0.536 (17.0)	0.566 (12.9)	0.346 (43.6)	0.204 (11.2)	0.350 (21.2)		0.204 (10.8)
Age	-0.641 (27.6)	-0.560 (10.0)	-0.636 (7.8)	-0.390 (39.7)	-0.268 (13.7)	-0.330 (14.4)	-0.232 (31.3)	-0.233 (12.6)
Married	-0.398 (29.0)	-0.449 (14.3)	-0.525 (12.1)	-0.495 (50.4)	-0.433 (33.7)	-0.469 (33.6)	-0.447 (56.9)	-0.429 (34.1)
Educated	0.0638 (4.1)	0.0647 (1.8)	0.0693 (1.4)	0.0772 (17.9)	-0.0983 (8.5)	-0.0865 (6.9)	0.0494 (12.4)	-0.0999 (7.5)
Female	-0.160 (5.3)	-0.350 (5.0)	-0.375 (3.9)	-0.0853 (8.3)	-0.340 (11.9)	-0.253 (9.8)	-0.268 (28.5)	-0.354 (11.9)
N Observations	81,285	16,896	8,855	1,961,708	268,300	217,225	1,910,441	258,102
Pseudo R2	0.1025	0.1533	0.1619	0.1110	0.1699	0.1513	0.2455	0.1789

Notes: Robust t-statistics in parentheses.